

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehponline@niehs.nih.gov](mailto:ehponline@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

# **Assessment of the Probability of Autochthonous Transmission of Chikungunya Virus in Canada under Recent and Projected Climate Change**

Victoria Ng, Aamir Fazil, Philippe Gachon, Guillaume Deuymes, Milka Radojević, Mariola Mascarenhas, Sophiya Garasia, Michael A. Johansson, and Nicholas H. Ogden

### **Table of Contents**

**Figure S1** Comparison between the bias-corrected data from CRCM5-CanESM2-RCP4.5 and CRCM5-CanESM2-RCP8.5 with data from other RCMs and their ensemble mean.

**Figure S2** Risk categories for autochthonous CHIKV transmission by *Ae. albopictus* in Canada derived from combining the climatic suitability for CHIKV transmission potential ( $R_0$ ) with the climatic suitability for the presence of *Ae. albopictus* (SIG index) using the 75<sup>th</sup> percentile value of  $R_0$  distributed across temperature range 10°C to 40°C as the cut-off.

**Figure S3** Risk maps for autochthonous CHIKV transmission in Canada based solely on CHIKV transmission potential ( $R_0$ ) using the 75<sup>th</sup> percentile value of  $R_0$  distributed across temperature range of 10°C to 40°C as the cut-off.

**Figure S4** Duration in months where mean  $R_0 > 1.0$  (mean monthly temperature between  $\geq 22.8^\circ\text{C}$  and  $33.6^\circ\text{C}$ ) in Canada based solely on CHIKV transmission potential ( $R_0$ ) using the 75<sup>th</sup> percentile value of  $R_0$  distributed across temperature range 10°C to 40°C as the cut-off.

**Figure S5** Risk maps for autochthonous CHIKV transmission in Canada combining the climatic suitability for CHIKV transmission potential ( $R_0$ ) with the climatic suitability for the presence of *Ae. albopictus* (SIG index) using the 75<sup>th</sup> percentile value of  $R_0$  distributed across temperature range 10°C to 40°C as the cut-off.

**Figure S6** Duration in months for potential autochthonous CHIKV transmission in Canada combining the climatic suitability for CHIKV transmission potential ( $R_0$ ) with the climatic suitability for the presence of *Ae. albopictus* (SIG index) using the 75<sup>th</sup> percentile value of  $R_0$  distributed across temperature range 10°C to 40°C as the cut-off.